









AcuPoint Analysis

DIGESTIVE DISTORDERS



- C12** 4 cun above the navel, along the mid-line.
- G05** On the spine between the first and second lumbar 1.5 cun above a the waist on a "normal-waisted" person.
- K20** The width of 5 thumbs above the navel and one thumb to either side of the abdominal mid-line.
- LI04** On the back of the hand between the thumb and index finger metacarpal bones. Find the spot midway on the bone that connects the index finger to the wrist and probe inward toward the main body of the hand.
- LV13** On the side of the abdomen at the free end of the 11th floating rib.
- P06** The width of two thumbs (2 Cun) above the most prominent crease on the inner wrist, in line with the middle finger.
- SP06** The width of one hand (3 thumbs) above the crown of the inner ankle, just behind the shin bone on the front of the leg.
- ST36** The width of one hand (3 Cun) below the bottom of the kneecap, and the width of one thumb (1 Cun) toward the outside of the leg (direction of the small toe.) Found in the trough or valley where the two most prominent bones of the calf come together.
- C06** 1.5 Cun below the navel.
- LV14** The space between the sixth and seventh ribs 3 cun directly below the nipples.
- UB21** 1.5 Cun from the spine midline, between the 12th thoracic vertebra and the 1st lumbar vertebra. 2.5 cun above the waist line. Easily stimulated with the Back-eze.
- UB25** 1.5 cun from the spine midline, on a level between the 4th and 5th lumbar vertebra. Just above an imaginary line drawn between the mid-forearms. Easily stimulated with the Back-eze.