
AcuPoint Analysis PARKINSON'S DISEASE 2

Points



AcuPoint Analysis PARKINSON'S DISEASE 2

(DESCRIPTIONS)

Point	Description
LI04	On the back of the hand between the thumb and index finger metacarpal bones. Find the spot midway on the bone that connects the index finger to the wrist and probe inward toward the main body of the hand.
SP06	The width of one hand (3 thumbs) above the crown of the inner ankle, just behind the shin bone on the front of the leg.
UB60	In the hollow (or valley) behind the crown of the outer ankle.
LI15	At the point of the shoulders. Raise your arm slightly above the level of your shoulder and place your Acu-Ki in the forward most dimple; then keeping your Acu-Ki in the dimple, lower your arm and begin stimulation.
SI08	At the elbow between the tip of the elbow and the bone tip of the upper arm on the inside of the arm.
H03	At the extreme end of the inner crease of the elbow. Bend the arm tightly and place the Acu-Ki at the extreme inner edge of the elbow crease formed. Relax your arm and begin the stimulation.
Т03	On the back of the hand behind the web of the fourth and the fifth finger and in a depression just behind the knuckle joints.
P03	In the deepest hollow of the inner elbow, along the crease, between the two ligaments.
P07	On the most prominent crease of the inner wrist, in line with the middle finger.
K03	The width of two thumbs above the crown of the inner ankle, still on the inner side of the leg, but almost to the rear of the leg. Do not strongly stimulate this point after the third month of pregnancy.
GB30	Near the "ball-joint" of the hips. Squeeze the buttocks together tightly, place the Acu-Ki in the depression formed at the hip, relax the buttocks and begin the stimulation.
LV03	Two cun above the separation between the biggest and second toes, on the top side of the foot.
G24	In the middle of the natural hairline, 3 cun above the center of the eyebrows.